



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

which permits two readings to be made with a very short time period between. An atmometer cup is mounted over a reservoir from which it may be cut off at will by means of a glass cock. It is also connected with a finely graduated burette from which the water will be drawn when the reservoir cock is closed. A reading can be made in a very short time at any desired intervals, and the average evaporating power for the period of observation can be calculated. Comparison of different environments is easily made.—CHARLES A. SHULL.

Embryo sac of *Oenothera*.—ISHIKAWA¹⁶ has published a very full account of the behavior of the gametophytes and the fertilization phenomena in *Oenothera nutans* and *O. pycnocarpa*, as well as in their hybrids. These two species were formerly included in *O. biennis*. The embryo sac arises from either the chalazal or micropylar megaspore, and often both develop simultaneously into complete embryo sacs. The sac is tetranucleate, lacking the antipodals and one of the polar nuclei. In fertilization one of the male nuclei fuses with the remaining polar nucleus, resulting in diploid endosperm. Self-sterility of some of the hybrids is due to feeble growth of the pollen tube. Tetranucleate embryo sacs occur also in *Ludwigia*, *Gaura*, *Godetia*, and *Circaea*.—J. M. C.

Iron in nutrient solutions.—CORSON and BAKKE,¹⁷ working upon wheat and Canada field peas, have studied the relative merits of ferrous and ferric phosphates in nutrient solution. They find that iron in the nutrient solution is more important than generally considered; that ferric phosphate is more effective than ferrous phosphate, especially for wheat; and that ferric phosphate in the concentration suggested by SHIVE (0.0044 grams per liter) gives maximum dry weight.—WM. CROCKER.

Polyembryony.—HARVEY,¹⁸ in connection with recording a case of polyembryony in *Quercus alba*, has given a summary of the recorded cases of polyembryony in angiosperms. The list includes 36 cases, scattered through "15 of the 49 alliances." In the case of *Quercus* reported two vigorous embryos occurred in the acorn, and it is of special interest because this is said to be the first reported case of polyembryony "in the first 13 alliances of the Archichlamydeae."—J. M. C.

¹⁶ ISHIKAWA, M., Studies on the embryo sac and fertilization in *Oenothera*. Ann. Botany 32:277-317. pl. 7. figs. 14. 1918.

¹⁷ CORSON, G. E., and BAKKE, A. L., The use of iron in nutrient solution for plants. Proc. Iowa Acad. Sci. 24:477-482. 1917.

¹⁸ HARVEY, LEROY H., Polyembryony in *Quercus alba*. Mich. Acad. Sci. Rep. 1917. 329-331.